

## Effect of spacing on growth and yield of fluted pumpkin, *Telfaria occidentalis* (Hook f.) in Okigwe, Nigeria

Onyekachi A. Uwalaka\*, Chukwudi E. Uwanaka, Hyginus N. Ibekwe, Blessing C. Usifo

National Horticultural Research Institute, P.M.B. 1076, Okigwe, Imo State, Nigeria

### ABSTRACT

A field experiment was conducted during the cropping season of 2017 to determine the most suitable plant spacing for optimum growth and yield of fluted pumpkin, *T. occidentalis* in Okigwe, Nigeria. The experiment was laid out in a Randomized Complete Block Design with three replications. The experimental treatments consisted of 40 cm × 40 cm (T<sub>1</sub>), 50 cm × 50 cm (T<sub>2</sub>), 60 cm × 60 cm (T<sub>3</sub>), and the standard spacing of 100 cm × 100 cm (T<sub>4</sub>) which served as control treatment. Data were collected on some growth and yield parameters at 2, 4, 6 and 8 weeks after planting (WAP). Analysis of variance was conducted on the data collected and significant means were separated using Fischer's Least Significant Difference (F-LSD) at 5 % significance level. The results revealed that plant spacing had no significant effect ( $P > 0.05$ ) on number of leaves, branches and vine length. Leaf area and the number of nodes per plant were significantly ( $P < 0.05$ ) influenced by plant spacing. Standard spacing of 100 cm × 100 cm (T<sub>4</sub>) and spacing of 60 cm × 60 cm (T<sub>3</sub>) ranked first and second with 10.86 and 9.70 nodes per plant, and 33.40 cm<sup>2</sup> and 33.03 cm<sup>2</sup> for leaf area respectively. Fresh and dry matter yields performed in the order T<sub>4</sub> > T<sub>3</sub> > T<sub>2</sub> > T<sub>1</sub>.

**Keywords:** Plant spacing, *Telfairia occidentalis*, vine length, yield

### INTRODUCTION

Fluted pumpkin, *Telfairia occidentalis* (Hook. f.) originated from West Africa's high – rainfall forest belt. The largest diversity in plant populations can currently be found in south-eastern Nigeria (Schippers, 2002). The leaves and seeds are very nutritious and frequently consumed by 35 million people, including in Western Cameroon and Benin, where the immature edible seeds are preferred rather than the leaves (Egbekan, *et al*, 1998). In Nigeria, the Igbos calls it “ugu” and the Yorubas use the name uguwu. In Cameroon, it is referred to as ekobon (Odiaka, 2001).

Plant spacing is an important agronomic practice which varies from one crop to another. It is a common practice by farmers to ensure good yield and harvest. Good spacing enhances plant development and growth, with improved consequential yield. Odiaka, (2001) and Odiaka *et al.* (2008) had documented the benefits of good spacing on crop yield. However, farmers in southeast Nigeria practice different plant spacing on *T. occidentalis* with varying results in terms of shoots, leaf,

and pod yield. This disparity necessitated the desire to conduct this experiment adopting different plant spacing methods in order to ascertain the most suitable plant spacing for fluted pumpkin, *T. occidentalis* in Okigwe agroecological zone of Nigeria for optimum growth and yield performance.

### MATERIALS AND METHODS

The experiment was conducted at the Vegetable Research Farm of NIHORT Mbato Outstation, Okigwe, Nigeria. The site was cleared manually with cutlasses, pulverized and planting beds of dimension 3m x 3m (9m<sup>2</sup>) were made with hand hoes. The experimental treatments consisted of 4 plant spacing dimensions; 40 cm × 40 cm (T<sub>1</sub>), 50 cm × 50 cm (T<sub>2</sub>), 60 cm × 60 cm (T<sub>3</sub>) and 100 cm × 100 cm (T<sub>4</sub>) laid out in Randomized Complete Block Design (RCBD) and replicated 3 times. The following agronomic parameters: number of leaves, vine length, collar diameter, number of branches,

\*Corresponding Author: Onyekachi A. Uwalaka, National Horticultural Research Institute, P.M.B. 1076, Okigwe, Imo State, Nigeria.

E-mail: [uwalaka.onyekashi@nihort.gov.ng](mailto:uwalaka.onyekashi@nihort.gov.ng)

Received: 27-10-2019

Revised: 12-11-2019

Published: 23-11-2019

number of nodes, and leaf area (cm<sup>2</sup>) were collected at 2, 4, 6 and 8 weeks after planting (WAP) using the method described by Obiefuna *et al.* (2012), while fresh and dry weights were collected at harvest from two (2) randomly selected plants. Data collected were subjected to Analysis of Variance (ANOVA) using Microsoft Excel Package 2007 Edition. The means were separated using Fisher's Least Significant Difference (F-LSD) at 5% significance level.

## RESULTS

### Effect of different plant spacing on the number of leaves and branches of *T. occidentalis*

The results presented in Tables 1 and 2 show the number of leaves and branches of *T. occidentalis* as influenced by plant spacing. No significant difference ( $P > 0.05$ ) existed among the treatment means. Results however showed that there was a steady increase in the number of leaves and branches as the plants grow. The highest number of leaves (21.29) was obtained from plants sown at 100 cm × 100 cm (T<sub>4</sub>) spacing, followed by T<sub>3</sub> (18.48) and T<sub>2</sub> (18.29) while the lowest number of leaves was recorded from T<sub>1</sub> (17.67). *T. occidentalis* in T<sub>4</sub> recorded the highest number of branches (2.60). This was closely followed by T<sub>3</sub> (2.24). However, no significant difference ( $P > 0.05$ ) was found between the treatment means for (T<sub>1</sub>) and (T<sub>2</sub>) plant spacing as both of them recorded the same number of branches per plant per plot respectively (Table 2).

### Effect of plant spacing on vine length

Main vine length and diameter of *T. occidentalis* are presented in Tables 3 and 4. Analysis of variance indicated that non-significant variation exists in vine length and diameter at 6 and 8 WAP. At 6 WAP, results showed that longest vine length was observed in T<sub>4</sub> (122.00 cm) while T<sub>3</sub> had 115.59 cm, followed by 108.06 cm recorded from T<sub>1</sub>, and T<sub>2</sub> (107.31 cm) in that order

respectively (Table 3). At 8 WAP, the vine lengths were 110.00, 130.43, 134.57 and 139.58 cm in T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub> respectively (Table 4) with the mean vine length ranging from 109.03 to 130.79 cm, and with T<sub>4</sub> spacing performing the best. The vine diameter at 6 WAP ranged from 0.30 to 0.34 cm, while at 8 WAP, the vine diameter ranged from 0.33 to 0.41 cm (Table 4). Similar trend was also observed on vine diameter with T<sub>4</sub> recording the highest vine diameter at 6 WAP (0.34 cm) and 8 WAP (0.41 cm) respectively. The order of performance regarding vine diameter was T<sub>4</sub> (0.38 cm) > T<sub>3</sub> (0.35 cm) > T<sub>2</sub> (0.34 cm) > T<sub>1</sub> (0.33 cm) [Table 4].

### Effect of spacing on the number of nodes and leaf area

Plant spacing had significant effect ( $P < 0.05$ ) on the mean number of nodes. The number of nodes per plant per plot was highest in T<sub>4</sub> (10.86), followed by T<sub>3</sub> (9.70), and then T<sub>1</sub> (8.1), while T<sub>2</sub> was found to have the least (7.8). The leaf area (cm<sup>2</sup>) per plant per plot was highest in T<sub>4</sub> (33.40cm<sup>2</sup>), and it then statistically significantly decreased progressively to 30.03, 28.86 and 17.94cm<sup>2</sup> in T<sub>3</sub>, T<sub>2</sub> and T<sub>1</sub> respectively (Table 5).

### Effect of plant spacing on the yield components of *T. occidentalis*

Fresh yield and dry matter yield of *T. occidentalis* as influenced by plant spacing are presented in Table 6. Analysis of variance conducted on the treatment means revealed that plant spacing did not have significant effect ( $P > 0.05$ ) on fresh and dry matter yields of *T. occidentalis*. Fresh yield was highest in T<sub>4</sub> (1148.88kg/ha), followed by T<sub>3</sub> (1137.5kg/ha), T<sub>2</sub> (1037 kg/ha) while the lowest fresh yield (1000.0 kg/ha) was obtained in T<sub>1</sub> (Table 6). Dry matter yield of *T. occidentalis* was highest (298.40 kg/ha) in *T. occidentalis* spaced at 100 cm x 100 cm (T<sub>4</sub>), T<sub>3</sub> closely ranked second in dry matter yield with 289.55 kg/ha, T<sub>2</sub> had 287.10 kg/ha, while T<sub>1</sub> had a dry matter yield of 262.50 kg/ha (Table 6).

**Table 1: Number of leaves of *T. occidentalis* as influenced by different plant spacing**

Plant spacing	Number of leaves				Mean
	2WAP	4WAP	6WAP	8WAP	
T1	7.99	15.00	21.70	26.00	17.67
T2	7.50	15.87	22.27	27.52	18.29
T3	8.00	14.96	21.24	29.72	18.48
T4	9.45	18.00	24.44	33.21	21.29
Mean	8.24	15.96	22.41	26.73	17.67
LSD 0.05	1.75	2.98	3.72	12.33	

WAP: Weeks after planting; T1: 40 x 40cm; T2: 50 x 50cm; T3: 60 x 60cm; T4: 100 x 100cm

**Table 2: Number of branches of *T. occidentalis* as influenced by different plant spacing**

Plant spacing	Number of branches				Mean
	2WAP	4WAP	6WAP	8WAP	
T1	0.9	1.33	2.04	3.37	1.97
T2	0.85	1.23	2.00	3.79	1.97
T3	1.3	1.92	2.29	3.43	2.24
T4	1.38	2.05	2.88	4.10	2.60
Mean	1.11	1.63	2.30	3.67	1.97
LSD 0.05	0.58	0.30	0.40	1.22	

WAP: Weeks after planting; T1: 40 x 40cm; T2: 50 x 50cm; T3: 60 x 60cm; T4: 100 x 100cm

**Table 3: Effect of plant spacing on main vine length**

Plant spacing	Main vine length (cm)		
	6WAP	8WAP	Mean
T1	108.06	110.00	109.03
T2	107.31	130.43	118.87
T3	115.59	134.57	125.08
T4	122.00	139.58	130.79
Mean	113.24	128.65	109.03
LSD 0.05	24.12	34.87	

WAP: Weeks after planting; T1: 40 x 40cm; T2: 50 x 50cm; T3: 60 x 60cm; T4: 100 x 100cm

**Table 4: Effect of plant spacing on diameter of vine length**

Plant spacing	Diameter of vine length (cm)		
	6WAP	8WAP	Mean
T1	0.32	0.33	0.33
T2	0.30	0.37	0.34
T3	0.33	0.37	0.35
T4	0.30	0.34	0.32
Mean	0.32	0.37	0.35
LSD 0.05	0.16	0.06	

WAP: Weeks after planting; T1: 40 x 40cm; T2: 50 x 50cm; T3: 60 x 60cm; T4: 100 x 100cm

**Table 5: Effect of plant spacing on the number of nodes and leaf area**

Plant spacing	Number of nodes per plant per plot	Leaf area (cm <sup>2</sup> ) per plant per plot
	T1	8.10
T2	7.80	28.86
T3	9.70	30.03
T4	10.86	33.40
Mean	9.12	27.56
LSD 0.05	1.33*	4.99*

\*Statistically significant; T1: 40 x 40cm; T2: 50 x 50cm; T3: 60 x 60cm; T4: 100 x 100cm

**Table 6: Cumulative yield components of *T. occidentalis* as affected by plant spacing**

Plant spacing	Yield (Kgha <sup>-1</sup> )	
	Fresh	Dry matter
T1	1000.00	262.50
T2	1037.50	287.10
T3	1137.50	289.55
T4	1148.88	298.40
Mean	1058.33	279.71
LSD 0.05	487.50	150.00

T1: 40 x 40cm; T2: 50 x 50cm; T3: 60 x 60cm; T4: 100 x 100cm

## DISCUSSION

Plant spacing improved the growth and yield components of *T. occidentalis* in Okigwe, Nigeria. Results affirmed the earlier report of Akanbi *et al.* (2006) that plant spacing is an important agronomic practice that enhances growth, vigor and the overall development of a crop. The results of this study also validates the earlier recommendation of 1m x 1m as standard plant spacing for *T. occidentalis* production and optimum yield by NIHORT Report, (2010). *T. occidentalis* planted at the spacing of 40 cm x 40 cm (T<sub>1</sub>) produced fewer fresh and dry matter yields. Ogbonna, (2009) who studied seed position in pod and effect on sex growth and yield of *T. occidentalis* reported no significant difference on harvestable/edible yields as influenced by sex ratio of *T. occidentalis*. It is a known fact that having more female plants on the field is considered more profitable since they also produce seeds. In the recent past, some efforts have been made by Anyim and Akoroda, (1983) to establish criteria for judicious selection of suitable planting distance for *T. occidentalis* cultivation as a way of enhancing the proportion of female plants in the field with the overall objective of improving the yield of the plant and better land area utilization. The performance of plants planted at 1m x 1m could be attributed to the fact that well-spaced plants had more space for growth and development, thus, the highest yields were recorded in plants spaced at T<sub>4</sub>.

## CONCLUSION

The results showed that *T. occidentalis* planted at 1m x 1m spacing had the highest mean number of leaves and branches as compared to other spacing used. This suggests that good planting distance should be used to obtain better yield in *T. occidentalis* production. However, the findings from this study showed that there was no significant difference ( $P>0.05$ ) in the number of branches and leaves per plant per plot in all the evaluated plant spacing. Whereas, the planting distance had no significant effect ( $P>0.05$ ) on the main vine lengths and collar diameter of *T. occidentalis*, both the fresh and dry matter yields were highest in plants spaced at T<sub>4</sub>, and with the order of performance being T<sub>4</sub>>T<sub>3</sub>>T<sub>2</sub>>T<sub>1</sub>.

## Source of support

Nil.

## Conflict of interest

None declared.

## REFERENCES

- Akanbi WB, Togun AO, Adediran JA, Ilupeju EAO (2010). Growth, dry matter and fruit yields components of okra under organic and inorganic sources of nutrients. *American-Eurasian Journal of Sustainable Agriculture* 4(1): 1-13.
- Anyim OA, Akoroda MO (1983). Determination of optimal sex-ratio for maximum fruit yield in *Telfairia occidentalis* - NIHORT Annual Report 1983. Ibadan, Nigeria. NIHORT.
- Egbekun MK, Nda-Suleiman EO, Akinyeye O (1998). Utilization of fluted pumpkin fruit (*Telfairia occidentalis*) in marmalade manufacturing. *Plant Foods Hum. Nutr.* 52(2): 171-6.
- NIHORT (2010). National Horticultural Research Institute Annual Report 2010. Ibadan, Nigeria: NIHORT.
- Obiefuna JC, Ibeawuchi II, Baiyeri KP, Obiefuna CA (2012). Field experimentation and communication for agricultural development, Owerri, Nigeria: DFC Publishers.
- Odiaka NI, Akoroda MO, Odiaka EC (2008). Diversity and production methods of fluted pumpkin (*Telfairia occidentalis* Hook F.); experience with vegetable farmers in Makurdi, Nigeria. *African J. Biotech.* 7(8): 944-954.
- Odiaka NI (2001). Survey on the production and supply of *Telfairia occidentalis* in Makurdi, Benue State. Crop Production Department, University of Agriculture.
- Ogbonna PE (2009). Pod portion and type effects on sex, growth and yield of fluted pumpkin. *Afr. Crop Sci. J.* 16(3): 185-190.
- Schippers RR (2000). African indigenous vegetables, an overview of the cultivated species. London, UK: University of Greenwich, Natural Resources Institute.

**How to cite this article:** Uwalaka OA, Uwanaka CE, Ibekwe HN, Usifo BC (2019). Effect of spacing on growth and yield of fluted pumpkin *Telfaria occidentalis* (Hook f.) in Okigwe, Nigeria. *Int. Arch. Multidis. Study* 1(1): 13-16.